



Soils Map



State: **Minnesota**
 County: **Kandiyohi**
 Location: **11-119N-36W**
 Township: **St. Johns**
 Acres: **112.27**
 Date: **6/17/2021**



©2021 AgriData, Inc.

Soils data provided by USDA and NRCS.

Area Symbol: MN067, Soil Area Version: 19

Code	Soil Description	Acres	Percent of field	PI Legend	Non-Irr Class *c	Productivity Index	Bromegrass alfalfa	Bromegrass alfalfa hay	Corn	Oats	Soybeans	Spring wheat	*n NCCPI Soybeans
817	Canisteo-Seaforth complex, 0 to 3 percent slopes	29.87	26.6%		IIw	94							83
981	Canisteo-Harps loams	26.06	23.2%		IIw	92	4.9	3.8	168	81	50	48	75
1900	Okoboji-Canisteo depressional complex, 0 to 1 percent slopes	11.30	10.1%		IIIw	86							74
423	Seaforth loam, 1 to 3 percent slopes	10.49	9.3%		IIIs	95							86
L163A	Okoboji silty clay loam, 0 to 1 percent slopes	9.76	8.7%		IIIw	86							74
897B	Seaforth-Swanlake loams, 2 to 6 percent slopes	6.24	5.6%		IIIs	90	5.4	4.2	165	85	50	52	71
927	Harps-Glencoe-Seaforth complex, 0 to 3 percent slopes	5.75	5.1%		IIw	90							82
810	Corriff-Fieldon complex	4.65	4.1%		IIw	79	5.1	3.8	145	82	43	49	60
L223B	Amiret-Swanlake loams, 2 to 6 percent slopes	4.48	4.0%		IIe	92							83
446B	Normania loam, 2 to 5 percent slopes	1.91	1.7%		IIe	94	5.8	4.5	172	87	52	52	67
999B	Ves-Swanlake-Hawick complex, 2 to 6 percent slopes	1.36	1.2%		IIe	74	5.1	3.9	135	80	41	48	52

140	Spicer silty clay loam, 0 to 2 percent slopes	0.40	0.4%		IIw	91							88
Weighted Average						90.7	1.8	1.4	58.7	29.4	17.5	17.5	*n 77.4

*n: The aggregation method is "Weighted Average using all components"

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.